

A VERTEX OPERATOR REPRESENTATION OF SOLUTIONS TO THE GUREVICH-ZYBIN HYDRODYNAMICAL EQUATION

YAREMA A. PRYKARPATSKY^{1,2}, DENIS BLACKMORE³, JOLANTA GOLENIA,
AND ANATOLIY K. PRYKARPATSKY^{5,6}

ABSTRACT. An approach based on the spectral and Lie - algebraic techniques for constructing vertex operator representation for solutions to a Riemann type hydrodynamical hierarchy is devised. A functional representation generating an infinite hierarchy of dispersive Lax type integrable flows is obtained.

1. INTRODUCTION

Nonlinear hydrodynamic equations are of constant interest still from classical works by B. Riemann, who had extensively studied them in general three-dimensional case, having paid special attention to their one-dimensional spatial reduction, for which he devised the generalized method of characteristics and Riemann invariants. These methods appeared to be very effective [1] in investigating many types of nonlinear spatially one-dimensional systems of hydrodynamical type and, in particular, the characteristics method in the form of a "reciprocal" transformation of variables has been used recently in studying a so called Gurevich-Zybin system [2, 3] in [8] and a Whitham type system in [6, 5]. Moreover, this method was further effectively applied to studying solutions to a generalized [5] (owing to D. Holm and M. Pavlov) Riemann type hydrodynamical system

$$(1.1) \quad D_t^N u = 0, \quad D_t := \partial/\partial t + u\partial/\partial x,$$

where $N \in \mathbb{Z}_+$ and $u \in C^\infty(\mathbb{R}^2; \mathbb{R})$ is a smooth function. Making use of novel methods, devised in [24, 7] and based both on the spectral theory [9, 16, 19, 18] and the differential algebra techniques, the Lax type representations for the cases $N = 1, 4$ were constructed in explicit form.

In this work we are interested in constructing a so called vertex operator representation [13, 14, 21, ?] for solutions to the Gurevich-Zybin hydrodynamical hierarchy (1.1) at $N = 2$:

$$(1.2) \quad \begin{cases} D_t u = u_t + uu_x = v, \\ D_t v = v_t + uv_x = 0, \end{cases}$$

making use an approach recently devised in [22, 23] for the case of the classical AKNS hierarchy of integrable flows, and which can be easily generalized for treating the problem for arbitrary integers $N \in \mathbb{Z}_+$.

2. A VERTEX OPERATOR ANALYSIS

We begin with a Lax type linear spectral problem [8, 5, 4] for the equation (1.1) at $N = 2$:

$$(2.1) \quad \begin{cases} D_t u = u_t + uu_x = v, \\ D_t v = v_t + uv_x = 0, \end{cases}$$

defined on the space of smooth real-valued 2π -periodic functions $(u, v)^\top \in M \subset C^\infty(\mathbb{R}/2\pi\mathbb{Z}; \mathbb{R}^2)$:

$$(2.2) \quad df/dx = \ell[u, v; \lambda]f, \quad \ell[u, v; \lambda] := \begin{pmatrix} -\lambda u_x/2 & -v_x \\ \lambda^2/2 & \lambda u_x/2 \end{pmatrix},$$

Date: present.

1991 *Mathematics Subject Classification.* Primary 58A30, 56B05 Secondary 34B15; PACS 02.30.Ik, 02.10.Ox, 47.35.Fg .

Key words and phrases. Lax type integrability, vertex operator representation, Lax integrability, Lie-algebraic approach.

where, by definition, $v := D_t u$, $f \in L_\infty(\mathbb{R}/2\pi\mathbb{Z}; \mathbb{C}^2)$ and $\lambda \in \mathbb{C}$ is a spectral parameter. Assume that a vector function $(u, v)^\top \in M$ depends parametrically on the infinite set $t := \{t_1, t_2, t_3, \dots\} \in \mathbb{R}^{\mathbb{Z}_+}$ in such a way that the generalized Floquet spectrum [9, 15, 18] $\sigma(\ell) := \{\lambda \in \mathbb{C} : \sup_{x \in \mathbb{R}} \|f(x; \lambda)\|_\infty < \infty\}$ of the linear problem (2.2) persists in being parametrically iso-spectral, that is $d\sigma(\ell)/dt_j = 0$ for all $t_j \in \mathbb{R}$. The iso-spectrality condition gives rise to a hierarchy of commuting to each other nonlinear bi-Hamiltonian dynamical systems on the functional manifold M in the general form

$$(2.3) \quad \frac{d}{dt_j}(u(t), v(t))^\top = -\vartheta \text{grad} H_j[u, v] := K_j[u(t), v(t)],$$

where $K_j : M \rightarrow T(M)$ and $H_j \in \mathcal{D}(M)$, $j \in \mathbb{Z}_+$, are, respectively, vector fields and conservation laws on the manifold M , which were before described in [5, 4, 7],

$$(2.4) \quad \vartheta := \begin{pmatrix} 0 & \partial \\ \partial & 0 \end{pmatrix}$$

is a Poisson structure on the manifold M and, by definition,

$$(2.5) \quad \begin{pmatrix} u(t) \\ v(t) \end{pmatrix} := \begin{pmatrix} u(x, t_1, t_2, t_3, \dots) \\ v(x, t_1, t_2, t_3, \dots) \end{pmatrix}$$

for $t \in \mathbb{R}^{\mathbb{N}}$.

It is well known [15, 18, 9, 16] that the Casimir invariants, determining conservation laws for dynamical systems (2.3), are generated by the suitably normalized monodromy matrix $\tilde{S}(x; \lambda) \in \text{End } \mathbb{C}^2$ of the linear problem (2.2)

$$(2.6) \quad \tilde{S}(x; \lambda) = k(\lambda)S(x; \lambda) - \frac{k(\lambda)}{2} \text{tr} S(x; \lambda),$$

where $F(y, x; \lambda) \in \text{End } \mathbb{C}^2$ is the matrix solution to the Cauchy problems

$$(2.7) \quad \frac{d}{dy} F(y, x; \lambda) = \ell(y; \lambda) F(y, x; \lambda), \quad F(y, x; \lambda)|_{y=x} = \mathbf{I},$$

for all $\lambda \in \mathbb{C}$ and $x, y \in \mathbb{R}$, where $\mathbf{I} \in \text{End } \mathbb{C}^2$ is the identity matrix, $S(x; \lambda) := F(x + 2\pi, x; \lambda)$ is the usual monodromy matrix for the equation (2.7). Here the parameter $k(\lambda) \in \mathbb{C}$ is invariant with respect to flows (2.3) and is chosen in such a way that the asymptotic condition

$$(2.8) \quad \tilde{S}(x; \lambda) \in \tilde{\mathcal{G}}_-$$

as $\lambda \rightarrow \infty$ holds for all $x \in \mathbb{R}$. Here $\tilde{\mathcal{G}}_- \subset \tilde{\mathcal{G}}$, where $\tilde{\mathcal{G}} := \tilde{\mathcal{G}}_+ \oplus \tilde{\mathcal{G}}_-$ is the natural splitting into two affine subalgebras of positive and negative λ -expansions of the centrally extended [15, 25] affine current $\mathfrak{sl}(2)$ -algebra $\hat{\mathcal{G}} := \tilde{\mathcal{G}} \oplus \mathbb{C}$:

$$(2.9) \quad \tilde{\mathcal{G}} := \{a = \sum_{j \in \mathbb{Z}, j \ll \infty} a^{(j)} \otimes \lambda^j : a^{(j)} \in C^\infty(\mathbb{R}/2\pi\mathbb{Z}; \mathfrak{sl}(2; \mathbb{C}))\}.$$

The latter is endowed with the Lie commutator

$$(2.10) \quad [(a_1, c_1), (a_2, c_2)] := ([a_1, a_2], \langle a_1, da_2/dx \rangle),$$

where the scalar product is defined as

$$(2.11) \quad \langle a_1, a_2 \rangle := \text{res}_{\lambda=\infty} \int_0^{2\pi} \text{tr}(a_1 a_2) dx$$

for any two elements $a_1, a_2 \in \tilde{\mathcal{G}}$ with "res" and "tr" being the usual residue and trace maps, respectively. As the spectrum $\sigma(\ell) \subset \mathbb{C}$ of the problem (2.2) is supposed to be parametrically independent, flows (2.3) are naturally associated with evolution equations

$$(2.12) \quad d\tilde{S}/dt_j = [(\lambda^{j+1}\tilde{S})_+, \tilde{S}]$$

for all $j \in \mathbb{R}$, which are generated by the set $I(\hat{\mathcal{G}}^*)$ of Casimir invariants of the coadjoint action of the current algebra $\hat{\mathcal{G}}$ on a given element $\ell(x; \lambda) \in \tilde{\mathcal{G}}^* \cong \tilde{\mathcal{G}}_+$ contained in the space of smooth functionals $\mathcal{D}(\hat{\mathcal{G}})$. In particular, a functional $\gamma(\lambda) \in I(\hat{\mathcal{G}})$ if and only if

$$(2.13) \quad [\tilde{S}(x; \lambda), \ell(x; \lambda)] + \frac{d}{dx} \tilde{S}(x; \lambda) = 0,$$

where the gradient $\tilde{S}(x; \lambda) := \text{grad}\gamma(\lambda)(\ell) \in \tilde{\mathcal{G}}_-$ is defined with respect to the scalar product (2.11) by means of the variation

$$(2.14) \quad \delta\gamma(\lambda) := \langle \text{grad}\gamma(\lambda)(\ell), \delta\ell \rangle.$$

To construct the solution to matrix equation (2.13), we find preliminary a partial solution $\tilde{F}(y, x; \lambda) \in \text{End } \mathbb{C}^2$, $x, y \in \mathbb{R}$, to equation (2.7) satisfying the asymptotic Cauchy data

$$(2.15) \quad \tilde{F}(y, x; \lambda)|_{y=x} = \mathbf{I} + O(1/\lambda)$$

as $\lambda \rightarrow \infty$. It is easy to check that

$$(2.16) \quad \tilde{F}(y, x; \lambda) = \begin{pmatrix} \tilde{e}_1(y, x; \lambda) & -\frac{\tilde{\beta}(y; \lambda)}{\lambda} \tilde{e}_2(y, x; \lambda) \\ -\frac{\lambda}{\tilde{\alpha}(y; \lambda)} \tilde{e}_1(y, x; \lambda) & \tilde{e}_2(y, x; \lambda) \end{pmatrix},$$

is an exact functional solution to (2.7) satisfying condition (2.15), where we have defined

$$(2.17) \quad \begin{aligned} \tilde{e}_1(y, x; \lambda) &:= \exp\left\{\frac{\lambda}{2}[u(x) - u(y)] + \lambda \int_x^y \tilde{\alpha} dv(s)\right\}, \\ \tilde{e}_2(y, x; \lambda) &:= \exp\left\{\frac{\lambda}{2}[u(y) - u(x)] - \frac{\lambda}{2} \int_x^y \tilde{\beta} ds\right\}, \end{aligned}$$

with the vector-functions $\alpha^\pm \in C^\infty(\mathbb{R}/2\pi\mathbb{Z}; \mathbb{R})$ satisfying the following determining functional relationships:

$$(2.18) \quad \begin{aligned} \tilde{\alpha} &= u_x + (u_x^2 - 2v_x + \xi \tilde{\alpha})^{1/2}, \\ \tilde{\beta} &= u_x - (u_x^2 - 2v_x + \xi \tilde{\beta})^{1/2}, \end{aligned}$$

as $\xi := 1/\lambda \rightarrow 0$ and existing when the condition $\varphi(x, t) := \sqrt{u_x^2 - 2v_x} \neq 0$ on the manifold M at $t = 0 \in \mathbb{R}^N$.

The fundamental matrix $F(y, x; \lambda) \in \text{End } \mathbb{C}^2$ can be represented for all $x, y \in \mathbb{R}$ in the form

$$(2.19) \quad F(y, x; \lambda) = \tilde{F}(y, x; \lambda) \tilde{F}^{-1}(x, x; \lambda).$$

Consequently, if one sets $y = x + 2\pi$ in this formula and defines the expression

$$(2.20) \quad k(\lambda) := \lambda^{-1}[\tilde{e}_1(x + 2\pi, x; \lambda) - \tilde{e}_2(x + 2\pi, x; \lambda)]^{-1},$$

it follows from (2.6), (2.16) and (2.19) that the exact functional matrix representation

$$(2.21) \quad \tilde{S}(x; \lambda) = \begin{pmatrix} \frac{[\tilde{\alpha}(x; \lambda) + \tilde{\beta}(x; \lambda)]}{2\lambda[\tilde{\alpha}(x; \lambda) - \tilde{\beta}(x; \lambda)]} & \frac{\tilde{\alpha}\tilde{\beta}}{\lambda^2[\tilde{\alpha}(x; \lambda) - \tilde{\beta}(x; \lambda)]} \\ -\frac{1}{[\tilde{\alpha}(x; \lambda) - \tilde{\beta}(x; \lambda)]} & \frac{[\tilde{\beta}(x; \lambda) + \tilde{\alpha}(x; \lambda)]}{2\lambda[\tilde{\beta}(x; \lambda) - \tilde{\alpha}(x; \lambda)]} \end{pmatrix},$$

satisfies the necessary condition (2.8) as $\lambda \rightarrow \infty$.

Remark 2.1. The invariance of the expression (2.20) with respect to the generating vector field (2.3) on the manifold M derives from the representation (2.19), the equations (2.13) and

$$(2.22) \quad \frac{d}{dt} \tilde{F}(y, x_0; \mu) = \frac{\lambda^3}{\mu - \lambda} \tilde{S}(x; \lambda) \tilde{F}(y, x_0; \mu),$$

which follows naturally from the determining matrix flows (2.12) upon applying the translation $y \rightarrow y + 2\pi$.

The matrix expression (2.21) gives rise to the following important functional relationships:

$$(2.23) \quad \frac{1 - \lambda(\tilde{s}_{11} - \tilde{s}_{22})}{2\tilde{s}_{21}} = \tilde{\alpha}, \quad \frac{-2\lambda^2 \tilde{s}_{12}}{1 - \lambda(\tilde{s}_{11} - \tilde{s}_{22})} = \tilde{\beta},$$

which allow to introduce in a natural way the vertex operator vector fields

$$(2.24) \quad X_\lambda^\pm = \exp(\pm D\lambda), \quad D_\lambda := \sum_{j \in \mathbb{Z}_+} \frac{1}{(j+1)} \lambda^{-(j+1)} \frac{d}{dt_{j+1}},$$

acting on an arbitrary smooth function $\eta \in C^\infty(\mathbb{R}^{\mathbb{Z}_+}; \mathbb{R})$ by means of the shifting mappings:

$$(2.25) \quad \begin{aligned} X_\lambda^\pm \eta(x, t_1, t_2, \dots, t_j, \dots) &:= \eta^\pm(x, t; \lambda) = \\ &= \eta(x, t_1 \pm 1/\lambda, t_2 \pm 1/(2\lambda^2), t_3 \pm 1/(3\lambda^3), \dots, t_j \pm 1/(j\lambda^j), \dots) \end{aligned}$$

as $\lambda \rightarrow \infty$. Namely, we following proposition holds.

Proposition 2.2. *The functional vertex operator expressions*

$$(2.26) \quad \begin{aligned} \tilde{\alpha}(x, t; \lambda) &= X_{\lambda}^{-} \alpha(x, t) = \alpha^{-}(x, t; \lambda), \\ \tilde{\beta}(x, t; \lambda) &= X_{\lambda}^{+} \beta(x, t) = \beta^{+}(x, t; \lambda) \end{aligned}$$

solve the functional equations (2.18), that is

$$(2.27) \quad \begin{aligned} \alpha^{-} &= u_x + (u_x^2 - 2v_x + \xi \alpha^{-})^{1/2}, \\ \beta^{+} &= u_x - (u_x^2 - 2v_x + \xi \beta^{+})^{1/2}, \end{aligned}$$

where $t \in \mathbb{R}^{\mathbb{Z}_+}$ and $\xi = 1/\lambda \rightarrow 0$.

Proof. To state this proposition it is enough to show that the following relationships hold:

$$(2.28) \quad \begin{aligned} \frac{d}{d\xi} \left[\frac{1 - \lambda(\tilde{s}_{11} - \tilde{s}_{22})}{2\tilde{s}_{21}} \right]_{\lambda=1/\xi} &= \frac{d}{dt} \left[\frac{1 - \lambda(\tilde{s}_{11} - \tilde{s}_{22})}{2\tilde{s}_{21}} \right]_{\lambda=1/\xi}, \\ \frac{d}{d\xi} \left[\frac{-8\lambda^2 \tilde{s}_{12}}{1 - \lambda(\tilde{s}_{11} - \tilde{s}_{22})} \right]_{\lambda=1/\xi} &= \frac{d}{dt} \left[\frac{-8\lambda^2 \tilde{s}_{12}}{1 - \lambda(\tilde{s}_{11} - \tilde{s}_{22})} \right]_{\lambda=1/\xi} \end{aligned}$$

for any parameter $\xi \rightarrow 0$, where by definition

$$(2.29) \quad \frac{d}{dt} := \frac{d}{d\xi} D_{\lambda} \Big|_{\lambda=1/\xi} = \sum_{j \in \mathbb{Z}_+} \xi^j \frac{d}{dt_{j+1}}$$

is a generating evolution vector field. Before doing this we find the evolution equation

$$(2.30) \quad \frac{d}{dt} \tilde{S}(x; \mu) = [\lambda^3 \frac{d}{d\lambda} \tilde{S}(x; \mu), \tilde{S}(x; \lambda)]$$

on the matrix $\tilde{S}(x; \mu)$ as $\mu, \lambda \rightarrow \infty$, which entails the following differential relationships:

$$(2.31) \quad \begin{aligned} d\tilde{s}_{11}/dt &= \lambda^3 (\tilde{s}_{21} d\tilde{s}_{12}/d\lambda - \tilde{s}_{12} d\tilde{s}_{21}/d\lambda), \\ d\tilde{s}_{22}/dt &= \lambda^3 (\tilde{s}_{12} d\tilde{s}_{21}/d\lambda - \tilde{s}_{21} d\tilde{s}_{12}/d\lambda), \\ d\tilde{s}_{22}/dt &= \lambda^3 [\tilde{s}_{12} \frac{d}{d\lambda} (\tilde{s}_{11} - \tilde{s}_{22}) - (\tilde{s}_{11} - \tilde{s}_{22}) \frac{d\tilde{s}_{12}}{d\lambda}], \\ d\tilde{s}_{11}/dt &= \lambda^3 [\tilde{s}_{21} \frac{d}{d\lambda} (\tilde{s}_{22} - \tilde{s}_{11}) - (\tilde{s}_{22} - \tilde{s}_{11}) \frac{d\tilde{s}_{21}}{d\lambda}]. \end{aligned}$$

Using these relationships (2.31), one can easily obtain by means of simple, but rather cumbersome calculations, the needed relationships (2.28). As their direct consequences the *vertex operator representations* (2.26) for the vector functions $\tilde{\alpha}, \tilde{\beta} \in C(\mathbb{R}^{\mathbb{Z}_+}; \mathbb{R})$ hold. $\square$

Now we take into account that, owing to the determining functional representations (2.18), that the limits ^{∞}

$$(2.32) \quad \begin{aligned} \lim_{\lambda \rightarrow \infty} \alpha^{-}(x, t; \lambda) &= u_x(x, t) + \varphi(x, t), \\ \lim_{\lambda \rightarrow \infty} \beta^{+}(x, t; \lambda) &= u_x(x, t) - \varphi(x, t), \quad \varphi(x, t) := \sqrt{u_x^2(x, t) - 2v_x(x, t)}, \end{aligned}$$

exist on the manifold M . Moreover, having iterated the functional relationships (2.18), one can find that

$$(2.33) \quad \begin{aligned} X_{\lambda}^{-} \alpha &= \alpha^{-} = u_x + \varphi + \xi \left(\frac{u_{xx}}{\varphi} + \frac{\varphi_x}{\varphi} \right) + \\ &+ \frac{\xi^2}{2} \left(\frac{u_{xx}^2 + 2u_{xx}\varphi_x - u_{3x}\varphi}{\varphi^3} + \frac{\varphi_{xx}\varphi + 5\varphi_x^2}{\varphi^3} \right) + O(\xi^3), \\ X_{\lambda}^{+} \beta &= \beta^{+} = u_x - \varphi - \xi \left(\frac{u_{xx}}{\varphi} - \frac{\varphi_x}{\varphi} \right) - \\ &- \frac{\xi^2}{2} \left(\frac{u_{xx}^2 - 2u_{xx}\varphi_x + u_{3x}\varphi}{\varphi^3} + \frac{\varphi_{xx}\varphi + 5\varphi_x^2}{\varphi^3} \right) + O(\xi^3), \end{aligned}$$

which immediately yield the higher Riemann type commuting nonlinear Lax integrable dispersive dynamical systems on the functional manifold M . For instance, making use of the relationships

$$(2.34) \quad \lim_{\lambda \rightarrow \infty} [\alpha^{-}(x, t; \lambda) \pm \beta^{+}(x, t; \lambda)]/2 = \begin{cases} u_x(x, t), \\ \varphi(x, t) \end{cases},$$

one easily obtains that

$$(2.35) \quad \frac{d}{dt_1} \begin{pmatrix} u_x \\ \varphi \end{pmatrix} = \begin{pmatrix} -u_{xx}/\varphi \\ -\varphi_x/\varphi \end{pmatrix}, \quad \frac{d}{dt_2} \begin{pmatrix} u_x \\ \varphi \end{pmatrix} = \begin{pmatrix} (u_{xx}^2 + 7\varphi_x^2)/\varphi^3 \\ (2u_{3x}\varphi - 4u_x\varphi_x)/\varphi^3 \end{pmatrix}, \dots,$$

and so on, where $\varphi = \sqrt{u_x^2 - 2v_x}$ and we took into account that the following asymptotic expansions hold

$$(2.36) \quad \begin{aligned} X_\lambda^- \alpha(x, t; \lambda) &= u_x + \varphi - \xi(u_{x,t_1} + \varphi_{t_1}) + \\ &+ \frac{\xi^2}{2}(u_{x,t_1,t_1} + \varphi_{t_1,t_1} - u_{x,t_2} - \varphi_{t_2}) + O(\xi^3), \\ X_\lambda^+ \beta(x, t; \lambda) &= u_x - \varphi + \xi(u_{x,t_1} - \varphi_{t_1}) + \\ &+ \frac{\xi^2}{2}(u_{x,t_1,t_1} - \varphi_{t_1,t_1} + u_{x,t_2} - \varphi_{t_2}) + O(\xi^3) \end{aligned}$$

as $\xi = 1/\lambda \rightarrow 0$.

It is worth here to mention that the scheme devised above for finding the corresponding vertex operator representations for the Riemann type equation (2.1) can be similarly generalized for treating others equations of the infinite hierarchy (1.1) when $N \geq 3$, having taking into account the existence of their suitable Lax type representations found before in recent works [24, 5, 4].

3. CONCLUDING REMARKS

The vertex operator functional representations of the solution to the Riemann type hydrodynamical equation (2.1) in the form (2.27) is crucially based on the representations (2.23) and evolution equations (2.28), which provide a very straightforward and transparent explanation of many of “miraculous” vertex operator calculations presented before both in [13, 14] and in [21]. It should be noted that the effectiveness of our approach to studying the vertex operator representation of the Riemann type hierarchy owes much to the important exact representation (2.21) for the corresponding monodromy matrix, whose properties are described by means of applying the standard [15, 18, 16, 20] Lie-algebraic techniques. As an indication of possible future research, it should also be mentioned that it would be interesting to generalize the vertex operator approach devised in this work to other linear spectral problems such as those related to dynamical systems with a parametrical spectral [11, 17, 18] dependence, spatially two-dimensional [10], Pavlov’s and heavenly [12] dynamical systems.

4. ACKNOWLEDGMENTS

D. Blackmore wishes to thank the National Science Foundation for support from NSF Grant CMMI - 1029809 and his coauthor for enlisting him in the efforts that produced this paper. A.K. Prykarpatsky cordially thanks Profs. N. Bogolubov (Jr.) for useful discussions of the results obtained.

REFERENCES

- [1] Whitham G.B. Linear and Nonlinear Waves . Wiley-Interscience, New York, 1974, 221p
- [2] Gurevich A.V. and Zybin K.P. Nondissipative gravitational turbulence. Sov. Phys.-JETP, 67 (1988), p.p. 1–12
- [3] Gurevich A.V. and Zybin K.P. Large-scale structure of the Universe. Analytic theory Sov. Phys. Usp. 38 (1995), p.p. 687–722
- [4] Golenia J., Bogolubov N. (jr.), Popowicz Z., Pavlov M. and Prykarpatsky A. A new Riemann type hydrodynamical hierarchy and its integrability analysis. *Preprint ICTP*, IC/2009/095, 2010
- [5] Golenia J., Pavlov M., Popowicz Z. and Prykarpatsky A. On a nonlocal Ostrovsky-Whitham type dynamical system, its Riemann type inhomogeneous regularizations and their integrability. SIGMA, 6, 2010, 1-13
- [6] Prykarpatsky A.K. and Prytula M.M. The gradient-holonomic integrability analysis of a Whitham-type nonlinear dynamical model for a relaxing medium with spatial memory. Nonlinearity 19 (2006) 2115–2122
- [7] Popowicz Z. and Prykarpatsky A. K. The non-polynomial conservation laws and integrability analysis of generalized Riemann type hydrodynamical equations. Nonlinearity 23 (2010) 2517–2537; arXiv:submit/0044844 [nlin.SI] 21 May 2010
- [8] Pavlov M. The Gurevich-Zybin system. J. Phys. A: Math. Gen. 38 (2005), p. 3823-384050
- [9] Novikov S.P. (Editor) Theory of Solitons, Moscow, Nauka Publ., 1980 (in Russian).
- [10] Zakharov V.E., Manakov S.M. Construction of multi-dimensional nonlinear integrable systems and their solutions. Funct. Anal. Appl., 1985, v. 19, N 2, p. 11-25 (in Russian)
- [11] Burtsev S.P., Zakharov V.E., Mikhaylov A.V. Inverse scattering problem with changing spectral parameter. Theor. Math. Phys., 1987, v. 70, N 3, p. 323-341 (in Russian)
- [12] Manakov S.M., Santini P.M. On the solutions of the second heavenly and Pavlov equations. J.Phys. A: Math and Theor., 2009, v.42, N 40, p. 404013-10
- [13] Newell A.C. Solitons in mathematics and physics. Arizona, SIAM Publ., 1985

- [14] Dickey L.A. Soliton equations and Hamiltonian systems. NY, World Scientific, 1991
- [15] Faddeev L.D., Takhtadjan L.A. Hamiltonian methods in the theory of solitons. NY, Springer, 1987
- [16] Prykarpatsky A. and Mykytyuk I. Algebraic integrability of nonlinear dynamical systems on manifolds: classical and quantum aspects. Kluwer Academic Publishers, the Netherlands, 1998, 553p.
- [17] Blackmore D., Prykarpatsky A.K., Samoilenko V.H. Nonlinear dynamical systems of mathematical physics. NY, World Scientific, 2011
- [18] Hentosh O.Ye., Prytula M.M. and Prykarpatsky A.K. Differential-geometric integrability fundamentals of nonlinear dynamical systems on functional manifolds. (The second revised edition), Lviv University Publisher, Lviv, Ukraine, 2006 (in Ukrainian)
- [19] Mitropolski Yu.A., Bogoliubov N.N. (Jr.), Prykarpatsky A.K., Samoilenko V.Hr. Integrable Dynamical Systems. Nauka dumka, Kiev, 1987 (in Russian)
- [20] Blaszkak M. Multi-Hamiltonian theory of dynamical systems. Springer, Berlin, 1998
- [21] Pritula G.M., Vekslerchik V.E. Conservation laws for the nonlinear Schrödinger equation in Miwa variables. Inverse Problems, 2002, 18, p. 1355 (arxiv:nlin.SI/0008034, 2000)
- [22] Blackmore D., Prykarpatsky A.K. and Prykarpatsky Y.A. Iso-spectrally integrable dynamical systems on discrete manifolds: analytical aspect. Opuscula Mathematica, 2011, N2.
- [23] The AKNS hierarchy revisited: A vertex operator approach and its Lie-algebraic structure. arXiv:1012.1024v1 [nlin.SI]
- [24] Prykarpatsky A.K., Artemovych O.D., Popowicz Z. and Pavlov M.V. Differential-algebraic integrability analysis of the generalized Riemann type and Korteweg–de Vries hydrodynamical equations. J. Phys. A: Math. Theor. 43 (2010) 295205 (13pp)
- [25] Reyman A.G., Semenov-Tian-Shansky. Integrable systems. Moscow-Izhevsk, R&C-Dynamics, 2003 (in Russian)

¹THE DEPARTMENT OF APPLIED MATHEMATICS AT THE UNIVERSITY, KRAKOW 30059, POLAND, AND,

²DEPARTMENT OF DIFFERENTIAL EQUATIONS OF THE INSTITUTE MATHEMATICS AT NAS, KYIV, UKRAINE
E-mail address: yarpry@gmail.com

³DEPARTMENT OF MATHEMATICAL SCIENCES AND CENTER FOR APPLIED MATHEMATICS AND STATISTICS, NEW JERSEY INSTITUTE OF TECHNOLOGY, NEWARK, NJ 07102, USA
E-mail address: deblac@m.njit.edu

⁴THE DEPARTMENT OF APPLIED MATHEMATICS AT THE AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, KRAKOW 30059, POLAND, GOLJOLS@TLEN.PL

⁵THE DEPARTMENT OF MINING GEODESY AND ENVIRONMENT ENGINEERING AT THE AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, KRAKOW 30059, POLAND, AND, ⁶DEPARTMENT OF ECONOMICAL CYBERNETICS AT THE IVAN FRANKO PEDAGOGICAL STATE UNIVERSITY, DROHOBYCH, LVIV REGION, UKRAINE
E-mail address: pryk.anat@ua.fm, prykanat@cybergal.com